

Shear viscosity of neutron matter from realistic nucleon-nucleon interactions

Omar Benhar^{1,2} and Marco Valli^{2,1}

¹*INFN, Sezione di Roma. I-00185 Roma, Italy*

²*Dipartimento di Fisica, Università "La Sapienza". I-00185 Roma, Italy*

(Dated: February 1, 2008)

The calculation of transport properties of Fermi liquids, based on the formalism developed by Abrikosov and Khalatnikov, requires the knowledge of the probability of collisions between quasi-particles in the vicinity of the Fermi surface. We have carried out a numerical study of the shear viscosity of pure neutron matter, whose value plays a pivotal role in determining the stability of rotating neutron stars, in which these processes are described using a state-of-the-art nucleon-nucleon potential model. Within our approach medium modifications of the scattering cross section are consistently taken into account, through an effective interaction obtained from the matrix elements of the bare interaction between correlated states. Inclusion of medium effects leads to a large increase of the viscosity at densities larger than $\sim 0.1 \text{ fm}^{-3}$.

PACS numbers: 21.65.+f, 26.60.+c, 97.60.Jd

Viscosity plays a pivotal role in determining the stability of rotating neutron stars. As Chandrasekhar first pointed out [1], emission of gravitational radiation (GR) following the excitation of non-radial oscillation modes may lead to the instability of rotating stars. While this effect would make all perfect fluid rotating stars unstable, in presence of viscosity dissipative effects damp the oscillations, and may prevent the onset of the instability. As a consequence, a quantitative understanding of the viscosity of neutron star matter is required to determine whether a mode is stable or unstable (for a recent review on neutron star oscillations and instabilities see, e.g., Ref. [2] and references therein).

Early estimates of the shear viscosity coefficients of neutron star matter were obtained in the 70s by Flowers and Itoh, who used the measured scattering phase shifts to estimate the neutron-neutron scattering probability [3, 4]. Based on these results, Cutler and Lindblom carried out a systematic study of the effect of the viscosity on neutron star oscillations, using a variety of different models of equation of state (EOS) of neutron star matter [5].

The procedure followed by the authors of Ref. [5], while allowing for a quantitative analysis of the damping of neutron star oscillations, cannot be regarded as fully consistent. Ideally, the calculation of transport properties of neutron star matter and the determination of its EOS should be carried out using the *same* dynamical model. The work discussed in this paper is aimed at making a first step towards this goal.

We have computed the shear viscosity of pure neutron matter using a realistic nucleon-nucleon (NN) potential, the Argonne v_{18} model [6], previously employed to obtain the state-of-the-art EOS of Akmal, Pandharipande and Ravenhall [7]. Within our approach, based on the formalism of Correlated-Basis-Function (CBF) perturbation theory [8, 9], medium modifications of the NN scattering cross section are also consistently taken into

account, through an effective interaction derived from the same NN potential.

The theoretical description of transport properties of normal Fermi liquids is based on Landau theory [10]. Working within this framework and including the leading term in the low-temperature expansion, Abrikosov and Khalatnikov [11] obtained the approximate expression of the shear viscosity coefficient

$$\eta_{AK} = \frac{1}{5} \rho m^* v_F^2 \tau \frac{2}{\pi^2 (1 - \lambda_\eta)}, \quad (1)$$

where ρ is the density, v_F is the Fermi velocity and m^* and τ denote the quasiparticle effective mass and lifetime, respectively. The latter can be written in terms of the angle-averaged scattering probability $\langle W \rangle$ according to

$$\tau T^2 = \frac{8\pi^4}{m^{*3}} \frac{1}{\langle W \rangle}, \quad (2)$$

with

$$\langle W \rangle = \int \frac{d\Omega}{2\pi} \frac{W(\theta, \phi)}{\cos \theta/2}. \quad (3)$$

Note that the scattering process involves quasiparticles on the Fermi surface. As a consequence, for any given density ρ , the scattering probability only depends on the angular variables θ and ϕ , the magnitude of all quasiparticle momenta being equal to the Fermi momentum $p_F = (3\pi^2 \rho)^{1/3}$. Finally, the quantity λ_η appearing in Eq.(1) is defined as

$$\lambda_\eta = \frac{\langle W(1 - 3 \sin^4 \theta/2 \sin^2 \phi) \rangle}{\langle W \rangle}. \quad (4)$$

The exact solution of the equation derived in Ref. [11], obtained by Brooker and Sykes [12, 13], reads

$$\eta = \eta_{AK} \frac{1 - \lambda_\eta}{4} \times \sum_{k=0}^{\infty} \frac{4k+3}{(k+1)(2k+1)[(k+1)(2k+1) - \lambda_\eta]}, \quad (5)$$

the size of the correction with respect to the result of Eq.(1) being $0.750 < (\eta/\eta_{AK}) < 0.925$.

Eqs.(1)-(5) show that the key element in the determination of the viscosity is the in-medium NN scattering cross section. In Ref. [14], the relation between NN scattering in vacuum and in nuclear matter has been analyzed under the assumption that the nuclear medium mainly affects the flux of incoming particles and the phase space available to the final state particles, while leaving the transition probability unchanged. Within this picture $W(\theta, \phi)$ can be extracted from the NN scattering cross section measured in free space, $(d\sigma/d\Omega)_{\text{vac}}$, according to

$$W(\theta, \phi) = \frac{16\pi^2}{m^{*2}} \left(\frac{d\sigma}{d\Omega} \right)_{\text{vac}} \quad (6)$$

where m^* is the nucleon effective mass and θ and ϕ are related to the kinematical variables in the center of mass frame through $E_{cm} = p_F^2(1 - \cos\theta)/(2m)$, $\theta_{cm} = \phi$.

The above procedure has been followed in Ref. [15], whose authors have used the available tables of vacuum cross sections obtained from partial wave analysis [16]. In order to compare with the results of Ref. [15], we have first carried out a calculation of the viscosity using Eqs.(1)-(6) and the free space neutron-neutron cross section obtained from the Argonne v_{18} potential

$$v_{ij} = \sum_{n=1}^{18} v_n(r_{ij}) O_{ij}^n. \quad (7)$$

In the above equation

$$O_{ij}^{n \leq 6} = [1, (\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j), S_{ij}] \otimes [1, (\boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j)] \quad (8)$$

where $\boldsymbol{\sigma}_i$ and $\boldsymbol{\tau}_i$ are Pauli matrices acting in spin and isospin space, respectively, and

$$S_{ij} = \frac{3}{r_{ij}^2} (\boldsymbol{\sigma}_i \cdot \mathbf{r}_{ij})(\boldsymbol{\sigma}_j \cdot \mathbf{r}_{ij}) - (\boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j). \quad (9)$$

The operators corresponding to $p = 7, \dots, 14$ are associated with the non static components of the NN interaction, while those corresponding to $p = 15, \dots, 18$ account for charge symmetry violations. Being fit to the full Nijmegen phase shifts data base, as well as to low energy scattering parameters and deuteron properties, the Argonne v_{18} potential provides an accurate description of the measured cross sections by construction.

In Fig. 1, we show the quantity ηT^2 as a function of density. Our results are represented by the solid line, while the dot-dash line corresponds to the results obtained from Eqs.(43) and (46) of Ref. [15] using the same effective masses, computed from the effective interaction discussed below. The differences between the two curves are likely to be ascribed to the correction factor of Eq.(5), not taken into account by the authors of Ref. [15], and to the extrapolation needed to determine the cross sections at small angles within their approach.

To gauge the model dependence of our results, we have replaced the full Argonne v_{18} potential with its simplified form, referred to as v'_8 [17], which only includes the six static operators of Eq.(8) and the two spin-orbit operators $\mathbf{L} \cdot \mathbf{S} \otimes [1, (\boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j)]$. This eight operators are the minimal set required to describe NN scattering in S and P states. The corresponding results, represented by the dashed line, show that using the v'_8 potential leads to a few percent change of ηT^2 over the density range corresponding to $1/4 < (\rho/\rho_0) < 2$, $\rho_0 = 0.16 \text{ fm}^{-3}$ being the equilibrium density of symmetric nuclear matter.




FIG. 1: (Color online) Neutron matter ηT^2 as a function of density. Solid line: results obtained from Eqs.(1)-(6) using the Argonne v_{18} potential and m^* computed from to the effective interaction described in the text. Dot-dash line: results obtained from Eqs.(43) and (46) of Ref. [15] using the same m^* . Dashed line: same as the solid line, but with the Argonne v_{18} replaced by its reduced form v'_8 .

To improve upon the approximation of Eq.(6) and include the effects of medium-modifications of the NN scattering amplitude, we have replaced the bare NN potential with an *effective interaction*, derived within the CBF approach as discussed in Ref. [18].

The *correlated* states of neutron matter are obtained from the Fermi gas (FG) states through the transformation

$$|n\rangle = F|n_{FG}\rangle, \quad (10)$$

where the operator F , embodying the correlation structure induced by the NN interaction, is written in the form

$$F = \mathcal{S} \prod_{ij} f_{ij}, \quad (11)$$

$\mathcal{S}$ being the symmetrization operator. The two-body correlation functions f_{ij} , whose operatorial structure reflects the complexity of the NN potential, can be written in the form

$$f_{ij} = \sum_{n=1}^6 f^n(r_{ij}) O_{ij}^n, \quad (12)$$

with the O_{ij}^n given by Eq.(8).

The effective interaction v_{eff} is defined by the relation

$$\frac{\langle n|H|n\rangle}{\langle n|n\rangle} = \langle n_{FG}|T + v_{\text{eff}}|n_{FG}\rangle, \quad (13)$$

where H is the full nuclear hamiltonian and T is the kinetic energy operator. Realistic models of H include, in addition to the NN potential v_{ij} , a three-nucleon potential V_{ijk} needed to account for the measured binding energies of the few-nucleon systems, as well as the empirical equilibrium properties of symmetric nuclear matter [19]. In this work, we follow the somewhat simplified approach originally proposed in Ref. [20], in which the main effect of the three-body force is taken into account through a density dependent modification of the intermediate range part of v_{ij} . Moreover, in view of the weak model dependence of ηT^2 (see Fig. 1), the full v_{18} potential is replaced by its reduced form v'_8 , and the contribution of the non static components is disregarded [18].

In order to obtain v_{eff} from Eq.(13) the expectation value of H in the correlated ground state is evaluated at the two-body level of the cluster expansion [18]. The resulting effective interaction reads

$$v_{\text{eff}} = \sum_{i<j} f_{ij}^\dagger \left[-\frac{1}{m}(\nabla^2 f_{ij}) - \frac{2}{m}(\nabla f_{ij}) \cdot \nabla + v_{ij} f_{ij} \right]. \quad (14)$$

The radial functions $f^n(r_{ij})$ of Eq.(12) are solutions of a set of Euler-Lagrange equations satisfying the boundary conditions $f^1(r_{ij} \geq d) = 1$, $f^n(r_{ij} \geq d) = 0$, for $n = 2, 3$ and 4, and $f^n(r_{ij} \geq d_t) = 0$, for $i = 5, 6$ (see, e.g., Ref. [20]).

The effective interaction of Eq.(14) has been tested computing the energy per particle of symmetric nuclear matter and pure neutron matter in first order perturbation theory using the FG basis. In Fig. 2 our results are compared to those of Refs. [7] and [21]. The calculations of Ref. [7] (solid lines) have been carried out using a variational approach based on the FHNC-SOC formalism, with a hamiltonian including the Argonne v_{18} NN potential and the Urbana IX three-body potential [19]. The results of Ref. [21] (dashed line of the lower panel) have been obtained using the v'_8 and the same three-body potential within the framework of the Auxiliary Field Diffusion Monte Carlo (AFDMC) approach. The results of Fig. 2 show that the effective interaction provides a fairly reasonable description of the EOS.

Note that our approach does not involve adjustable parameters. The correlation ranges d and d_t have been taken from Ref. [22], while the parameters entering the definition of the three-nucleon interaction (TNI) have been determined by the authors of Ref. [20] through a fit of nuclear matter equilibrium properties.




FIG. 2: (Color online) Energy per particle of symmetric nuclear matter (upper panel) and pure neutron matter (lower panel). The diamonds represent the results obtained using the effective interaction discussed in the text in first order perturbation theory with the FG basis, whereas the solid lines correspond to the results of Akmal Pandharipande and Ravenhall [7]. The dashed line of the lower panel represents the results of the AFDMC approach or Ref. [21].

Knowing the effective interaction, the in medium scattering probability can be readily obtained from Fermi's golden rule. The corresponding cross section at momentum transfer $\mathbf{q}$ reads

$$\frac{d\sigma}{d\Omega} = \frac{m^{*2}}{16\pi^2} |\hat{v}_{eff}(\mathbf{q})|^2, \quad (15)$$

$\hat{v}_{eff}$ being the Fourier transform of the effective potential. The effective mass can also be extracted from the quasiparticle energies computed in Hartree-Fock approximation.

In Fig. 3 the in-medium neutron-neutron cross section at $E_{cm} = 100$ MeV obtained from the effective potential, with $\rho = \rho_0$ and $\rho_0/2$, is compared to the corresponding free space result. As expected, screening of the bare interaction leads to an appreciable suppression of the scattering cross section.

Replacing the cross section in vacuum with the one defined in Eq.(15), the medium modified scattering probability can be readily obtained from Eq.(6). The resulting $W(\theta, \phi)$ can then be used to calculate ηT^2 from Eqs.(1)-(5).

The effect of using the medium modified cross section is illustrated in Fig. 4. Comparison between the solid and dashed lines shows that inclusion of medium modifications leads to a large increase of the viscosity, ranging




FIG. 3: (Color online) Differential neutron-neutron scattering cross section at $E_{cm} = 100$ MeV, as a function of the scattering angle in the center of mass frame. Solid line: cross section in vacuum, calculated with the v'_8 potential. Dot-dash line: medium modified cross section obtained from the effective interaction described in the text at $\rho = 0.08 \text{ fm}^{-3}$. Dashed line: same as the dot-dash line, but for $\rho = 0.16 \text{ fm}^{-3}$.

between $\sim 75\%$ at half nuclear matter density to a factor of ~ 6 at $\rho = 2\rho_0$. Such an increase is likely to produce appreciable effects on the damping of neutron star oscillations.




FIG. 4: (Color online) Solid line: density dependence of ηT^2 computed using the effective interaction described in the text. Dashed line: ηT^2 obtained from the free space cross section corresponding to the v'_8 potential.

In conclusion, we have computed the shear viscosity of pure neutron matter using an effective interaction derived from a dynamical model that can also be used to obtain the EOS. While our results are interesting in their own right, as they can be employed in a quantitative analysis

of the effect of viscosity on neutron stars oscillations, we emphasize that the work described in this paper should be seen as a first step towards the development of a general approach, allowing for a consistent calculation of the properties of neutron star matter.

The authors are grateful to V. Ferrari, for drawing their attention to the subject of this paper, and to R. Schiavilla, for providing a code for the calculation of the NN scattering cross section. Useful discussions with I. Bombaci are also gratefully acknowledged.

-
- [1] S. Chandrasekhar, Phys. Rev. Lett. **24**, 611 (1970).
 - [2] L. Lindblom, in *Gravitational Waves: a Challenge to Theoretical Astrophysics*, edited by V. Ferrari, J.C. Miller and L. Rezzolla, ICTP Lecture Notes Series Vol. 3 (ICTP, Trieste, 2001); http://users.ictp.it/~pub_off/lectures/vol3.html.
 - [3] E. Flowers and N. Itoh, Astrophys. J. **206**, 218 (1976).
 - [4] E. Flowers and N. Itoh, Astrophys. J. **230**, 847 (1979).
 - [5] C. Cutler and L. Lindblom, Astrophys. J. **314**, 234 (1987).
 - [6] R.B. Wiringa, V.G.J. Stoks and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
 - [7] A. Akmal, V.R. Pandharipande and D.G. Ravenhall, Phys. Rev. C **58**, 1804 (1998).
 - [8] J.W. Clark, Prog. Part. Nucl. Phys. **2**, 89 (1979).
 - [9] S. Fantoni and V.R. Pandharipande, Phys. Rev. C **37**, 1697 (1988).
 - [10] G. Baym and C. Pethick, *Landau Fermi-Liquid Theory* (John Wiley & Sons, New York, 1991).
 - [11] A.A. Abrikosov and I.M. Khalatnikov, Soviet Phys. JETP **5**, 887 (1957); Rep. Prog. Phys. **22**, 329 (1959).
 - [12] G.A. Brooker and J. Sykes, Phys. Rev. Lett. **21**, 279 (1968).
 - [13] J. Sykes and G.A. Brooker, Ann. Phys. (N. Y.) **56**, 1 (1970).
 - [14] V.R. Pandharipande and S.C. Pieper, Phys. Rev. C **45**, 791 (1992).
 - [15] D.A. Baiko and P. Haensel, Acta Phys. Pol. **30**, 1097 (1999).
 - [16] R.A. Arndt, L.D. Roper, R.A. Bryan, R.B. Clark, B.J. VerWest and P. Signell, Phys. Rev. D **28**, 97 (1983).
 - [17] B.S. Pudliner, V.R. Pandharipande, J. Carlson, S.C. Pieper and R.B. Wiringa, Phys. Rev. C **56**, 1720 (1997).
 - [18] S. Cowell and V.R. Pandharipande, Phys. Rev. C **73**, 025801 (2006).
 - [19] B.S. Pudliner, V.R. Pandharipande, J. Carlson and R.B. Wiringa, Phys. Rev. Lett. **74**, 4396 (1995).
 - [20] I. Lagaris and V.R. Pandharipande, Nucl. Phys. **359**, 349 (1981).
 - [21] A. Sarsa, S. Fantoni, K.E. Schmidt and F. Pederiva, Phys. Rev. C **68**, 024308 (2003).
 - [22] A. Akmal and V.R. Pandharipande, Phys. Rev. C **56**, 2261 (1997).